

Algorithm description document for retrieval of SCIAMACHY vertical tropospheric NO₂ columns

Akhila Padmanabhan¹
Randall V. Martin^{1,2,3}
Aaron van Donkelaar²
Christopher E. Sioris⁴
Kelly V. Chance³
Thomas P. Kurosu³

¹*Department of Chemistry, Dalhousie University, Halifax, NS, Canada*

²*Department of Physics and Atmospheric Sciences, Dalhousie University, Halifax, NS, Canada*

³*Harvard-Smithsonian Centre for Astrophysics, Cambridge, MA, USA*

⁴*Experimental Studies (ARQX), Air Quality Research Division, Environment Canada, Downsview, Ontario, Canada*

1. Introduction

Nitrogen Oxides (NO + NO₂), or NO_x, play an important role in climate change and air quality. NO_x largely controls tropospheric ozone production. NO_x oxidizes to nitric acid which impacts cloud properties and aerosol formation. Information from satellites provide a ‘top-down’ constraint for NO_x emissions.

2. Forward Model and Inversion Procedure

Following *Chance* [1998] and *Martin et al* [2002], we determine total slant columns of NO₂ by directly fitting backscattered radiance spectra from SCIAMACHY within the wavelength range of 429-452 nm, using measured reference spectra as described in section 3.

We remove the stratospheric column by first assuming that NO₂ over the central Pacific is primarily stratospheric and then subtracting the corresponding columns from the SCIAMACHY observations for the appropriate latitude and month. The resulting tropospheric columns are then corrected for the small amount of NO₂ over the Pacific using NO₂ columns obtained from GEOS-Chem simulations [*Martin et al.*, 2006, www.geos-chem.org].

An air mass factor calculation is used to convert the tropospheric slant columns into vertical columns. This formula for the AMF follows *Palmer et al.* [2001]:

$$AMF = AMF_G \int_0^{\infty} w(z)S(z)dz \quad (1)$$

where AMF_G is the geometric AMF, w is the scattering weight, and S is the shape factor. The geometric AMF takes into account the geometry of the sun, satellite and earth and is a function of the solar zenith angle and satellite viewing angle. The scattering weights refer to the instrument’s sensitivity to the trace gas as a function of altitude. We use the Linearized Discrete

Ordinate Radiative Transfer (LIDORT) model to calculate the scattering weights [Spurr, 2002]. The shape factor refers to the relative vertical distribution of NO₂ in the troposphere and is obtained from the GEOS-Chem chemical transport model. Monthly varying surface reflectivity fields are from Kleipool *et al.* [2008]. The aerosol correction uses daily local aerosol profiles obtained from a GEOS-Chem simulation of that scene [Martin *et al.*, 2003]. The AMF cloud correction depends on cloud fraction (what percentage of the observed scene is cloud-free), optical depth, and cloud height. The Fast Retrieval Scheme for Cloud Observables (FRESCO+) algorithm is used to correct for clouds [Wang *et al.*, 2008]. The vertical tropospheric NO₂ columns are obtained using the following calculation:

$$\text{tropospheric vertical column} = \frac{\text{tropospheric slant column}}{\text{AMF}}$$

3. Auxiliary Data needed

Spectral Fitting:	
NO ₂ at 243K	<i>Bogumil et al., 2003</i>
ozone at 223K	<i>Bogumil et al., 2003</i>
H ₂ O at 296K	<i>Rothman et al., 1998</i>
O ₂ -O ₂ at 296K	<i>Rothman et al., 1998</i>
Ring effect	<i>Chance and Spurr, 1997</i>
solar reference spectrum	<i>Chance and Kurucz, 2010</i>
AMF Calculation:	
cloud-top pressure	<i>Wang et al., 2008</i>
cloud fraction	<i>Wang et al., 2008</i>
surface reflectivity fields	<i>Kleipool et al. 2008</i>
NO ₂ shape profiles	<i>GEOS-Chem</i>
aerosol correction	<i>GEOS-Chem</i>

4. Sensitivity and Error Analysis

The typical 1-sigma uncertainty for each SCIAMACHY measurement is estimated to be 40% due to the AMF plus 1×10^{15} molecules cm⁻² caused by spectral fitting and subtraction of the stratospheric NO₂ column. The estimate for monthly mean uncertainty is $\pm(5 \times 10^{14}$ molecules cm⁻² + 30%) which could include systematic errors [Martin *et al.*, 2006].

5. Algorithm Validation

As described in Martin *et al.* [2006], the tropospheric NO₂ columns were validated with coincident *in situ* measurements taken from the NASA DC-8 and the NOAA WP-3D aircraft as part of the ICARTT campaign. The two measurement techniques were consistent within their uncertainty. The retrieved columns tended to be lower than the columns derived from the *in situ*

measurements in regions of enhanced NO₂ and higher in remote areas. The *in situ* measurements were also used to evaluate the GEOS-Chem simulated NO₂ vertical profiles used in the air mass factor formulation. Additionally, the GEOS-Chem vertical profiles were evaluated over the southeastern United States using aircraft measurements from TexAQS and SOS campaigns. Comparison of the AMFs calculated from the model profiles to those obtained from the *in situ* profiles generally showed agreement within a few percent [Martin *et al.*, 2004].

6. Recommendations for product validation

Martin et al. [2004] contains recommendations on the use of *in situ* measurements complemented with ground based observations. Vertically resolved measurements are needed that span the entire troposphere and are coincident in time and space with satellite overpasses. Validation should extend over large spatial regions over extended periods of time.

References:

Bogumil, K., et al. (2003), Measurements of molecular absorption spectra with the SCIAMACHY pre-flight model: Instrument characterization and reference data for atmospheric remote-sensing in the 230– 2380 nm region, *J. Photochem. Photobiol. A*, *157*, 167– 184.

Chance, K. (1998), Analysis of BrO measurements from the Global Ozone Monitoring Experiment, *Geophys. Res. Lett.*, *25*, 3335– 3338.

Chance, K. and R.L. Kurucz (2010) , An improved high-resolution solar reference spectrum for Earth's atmosphere measurements in the ultraviolet, visible, and near infrared, *JQSRT*, *111*, 1289-1295.

Chance, K. V., and R. J. D. Spurr (1997), Ring effect studies: Rayleigh scattering, including molecular parameters for rotational Raman scattering, and the Fraunhofer spectrum, *Appl. Opt.*, *36*, 5224– 5230

Kleipool, Q. L., M. R. Dobber, J. F. de Haan, and P. F. Levelt (2008), Earth surface reflectance climatology from 3 years of OMI data, *Journal of Geophysical Research-Atmospheres*, *113*(D18), doi: 10.1029/2008JD010290 ER.

Martin, R. V., K. Chance, D.J. Jacob, T.P. Kurosu, R.J.D. Spurr, E. Bucsela, J.F. Gleason, P.I. Palmer, I. Bey, A.M. Fiore, Q. Li, R.M. Yantosca, and R.B.A. Koelemeijer. (2002), An improved retrieval of tropospheric nitrogen dioxide from GOME, *J. Geophys. Res.*, *107*(D20), 4437, doi:10.1029/ 2001JD001027.

Martin, R. V., D. J. Jacob, K. Chance, T. P. Kurosu, P. I. Palmer, and M. J. Evans (2003), Global inventory of nitrogen oxide emissions constrained by space-based observations of NO₂ columns, *J. Geophys. Res.*, *108*(D17), 4537, doi:10.1029/2003JD003453.

Martin, R. V., D.D. Parrish, T.B. Ryerson, D.K. Nicks Jr., K. Chance, T.P. Kurosu, A. Fried, B.P. Wert, D.J. Jacob, and E. D. Sturges (2004), Evaluation of GOME satellite measurements of tropospheric NO₂ and HCHO using regional data from aircraft campaigns in the southeastern United States, *J. Geophys. Res.*, *109*, D24307, doi:10.1029/2004JD004869.

Martin, R.V., C.E. Sioris, K. Chance, T.B. Ryerson, T.H. Bertram, P.J. Wooldridge, R.C. Cohen, J.A. Neuman, A. Swanson, and F.M. Flocke (2006) Evaluation of space-based constraints on global nitrogen oxide emissions with regional aircraft measurements over and downwind of eastern North America, *J. Geophys. Res.*, *111*, D15308, doi:10.1029/2005JD006680.

Palmer, P. I., et al. (2001), Air mass factor formulation for spectroscopic measurements from satellites: Application to formaldehyde retrievals from the Global Ozone Monitoring Experiment, *J. Geophys. Res.*, *106*, 14,539–14,550.

Rothman, L. S., et al. (1998), The HITRAN molecular spectroscopic database and HAWKS (HITRAN atmospheric workstation): 1996 edition, *J. Quant. Spectrosc. Radiat. Transfer*, *60*, 665– 710.

Spurr, R.J.D. (2002), Simultaneous derivation of intensities and weighting functions in a general pseudo-spherical discrete ordinate radiative transfer treatment, *J. Quant. Spectrosc. Radiat. Transfer*, 75, 129-175.

Wang, P., P. Stammes, R. van der A, G. Pinardi, M. van Roozendaal (2008) FRESCO+: an improved O₂-A band cloud retrieval algorithm for tropospheric trace gas retrievals, *ACP*, 8, 6565-6576